

STEP Support Programme

STEP 3 Polar Coordinates and other Coordinate Geometry: Questions

1 2009 S3 Q1

The points S , T , U and V have coordinates (s, ms) , (t, mt) , (u, nu) and (v, nv) , respectively. The lines SV and UT meet the line $y = 0$ at the points with coordinates $(p, 0)$ and $(q, 0)$, respectively. Show that

$$p = \frac{(m - n)sv}{ms - nv},$$

and write down a similar expression for q .

Given that S and T lie on the circle $x^2 + (y - c)^2 = r^2$, find a quadratic equation satisfied by s and by t , and hence determine st and $s + t$ in terms of m , c and r .

Given that S , T , U and V lie on the above circle, show that $p + q = 0$.

2 2004 S3 Q4

The triangle OAB is isosceles, with $OA = OB$ and angle $AOB = 2\alpha$ where $0 < \alpha < \frac{\pi}{2}$. The semi-circle C_0 has its centre at the midpoint of the base AB of the triangle, and the sides OA and OB of the triangle are both tangent to the semi-circle. C_1, C_2, C_3, \dots are circles such that C_n is tangent to C_{n-1} and to sides OA and OB of the triangle.


Let r_n be the radius of C_n . Show that

$$\frac{r_{n+1}}{r_n} = \frac{1 - \sin \alpha}{1 + \sin \alpha}.$$

Let S be the total area of the semi-circle C_0 and the circles C_1, C_2, C_3, \dots . Show that

$$S = \frac{1 + \sin^2 \alpha}{4 \sin \alpha} \pi r_0^2.$$

Show that there are values of α for which S is more than four fifths of the area of triangle OAB .



3 1993 S3 Q2

The curve C has the equation $x^3 + y^3 = 3xy$.

- (i) Show that there is no point of inflection on C . You may assume that the origin is not a point of inflection.
- (ii) The part of C which lies in the first quadrant is a closed loop touching the axes at the origin. By converting to polar coordinates, or otherwise, evaluate the area of this loop.

4 2008 S3 Q3


The point $P(a \cos \theta, b \sin \theta)$, where $a > b > 0$, lies on the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

The point $S(-ea, 0)$, where $b^2 = a^2(1 - e^2)$, is a focus of the ellipse. The point N is the foot of the perpendicular from the origin, O , to the tangent to the ellipse at P . The lines SP and ON intersect at T . Show that the y -coordinate of T is

$$\frac{b \sin \theta}{1 + e \cos \theta}.$$

Show that T lies on the circle with centre S and radius a .



5 2011 S3 Q5


A movable point P has cartesian coordinates (x, y) , where x and y are functions of t . The polar coordinates of P with respect to the origin O are r and θ . Starting with the expression

$$\frac{1}{2} \int r^2 d\theta$$

for the area swept out by OP , obtain the equivalent expression

$$\frac{1}{2} \int \left(x \frac{dy}{dt} - y \frac{dx}{dt} \right) dt. \tag{*}$$

The ends of a thin straight rod AB lie on a closed convex curve C . The point P on the rod is a fixed distance a from A and a fixed distance b from B . The angle between AB and the positive x direction is t . As A and B move anticlockwise round C , the angle t increases from 0 to 2π and P traces a closed convex curve D inside C , with the origin O lying inside D , as shown in the diagram.



Let (x, y) be the coordinates of P . Write down the coordinates of A and B in terms of a, b, x, y and t .

The areas swept out by OA, OB and OP are denoted by $[A], [B]$ and $[P]$, respectively. Show, using $(*)$, that

$$[A] = [P] + \pi a^2 - af$$

where

$$f = \frac{1}{2} \int_0^{2\pi} \left(\left(x + \frac{dy}{dt} \right) \cos t + \left(y - \frac{dx}{dt} \right) \sin t \right) dt.$$

Obtain a corresponding expression for $[B]$ involving b . Hence show that the area between the curves C and D is πab .

